

ON MULTIDEGREES OF POLYNOMIAL AUTOMORPHISMS OF $\mathbb{C}^3$

JAKUB ZYGADŁO

ABSTRACT. We prove that for every pair of positive integers a, b there exists a number c_0 such that for every $c \geq c_0$ one can find a tame polynomial automorphism of $\mathbb{C}^3$ with multidegree equal to (a, b, c) .

Let $F = (F_1, \dots, F_n)$ be a polynomial automorphism of $\mathbb{C}^n$ (although the results in this paper are valid over an algebraically closed field k of characteristic 0, see for ex. [1] for general information about polynomial automorphisms). We define the multidegree of F as $\text{mdeg } F := (\deg F_1, \dots, \deg F_n)$, where $\deg$ means the total degree of a polynomial. It is an open question whether every n -tuple $(d_1, \dots, d_n)$ of positive integers is a multidegree of some polynomial automorphism of $\mathbb{C}^n$. The case $n = 1$ is obvious, since $\deg F = 1$ if F is a polynomial automorphism. In the case $n = 2$, due to Jung-van der Kulk theorem (cf. [1] and [3] for the original proof), we must have $\deg F_1 \mid \deg F_2$ or $\deg F_2 \mid \deg F_1$ for $F = (F_1, F_2)$ to be a polynomial automorphism of $\mathbb{C}^2$. Provided that $d_1 \mid d_2$ (or $d_2 \mid d_1$) one can also easily find a polynomial automorphism $F = (F_1, F_2)$ such that $\text{mdeg } F = (d_1, d_2)$; for example $F = (x, y + x^{d_2/d_1}) \circ (x + y^{d_1}, y)$. The case $n = 3$, however, seems much more complicated: one cannot even say (to the author's knowledge) whether there exists a polynomial automorphism with multidegree equal to $(3, 4, 5)!$ A recent result by M. Karaś ([2]) shows that if such an automorphism exists, it is not tame (see definition below and cf. [4]).

From now on we will focus our attention on the case $n = 3$ and write (a, b, c) instead of (d_1, d_2, d_3) for the given multidegree. Without loss of generality we may assume that $c \geq b \geq a$. We will calculate an integer c_0 (depending on a and b) such that for every $c \geq c_0$ one can find a polynomial automorphism of $\mathbb{C}^3$ with multidegree (a, b, c) . Moreover, all automorphisms we introduce are tame - they are compositions of linear and elementary maps ($F = (F_1, \dots, F_n)$ is elementary if for some $j \leq n$ and a polynomial g in $n - 1$ variables $F_j = X_j + g(X_1, \dots, \widehat{X_j}, \dots, X_n)$ and $F_i = X_i$ for $i \neq j$). First, let us prove some simple facts about multidegrees.

Fact 1. *Let $c \geq b \geq a$ and*

- (1) *$c = ka + lb$ for some $k, l \geq 0$ or*

Date: March 31, 2009.

2000 Mathematics Subject Classification. 14R10.

- (2) $a|b$ or
- (3) $a \leq 2$.

Then there exists a tame automorphism of $\mathbb{C}^3$ with multidegree (a, b, c) .

Proof. We will explicitly show the required automorphism:

- (1) One can take $F = (x, y, z + x^k y^l) \circ (x + z^a, y + z^b, z)$.
- (2) Let $b = da$ and take $F = (x, y + x^d, z) \circ (x + y^a, y, z + y^c)$.
- (3) Case $a = 1$ is trivial, so let $a = 2$. If $2|b$ we are done due to (2). So let b - odd and $c = b + m$, $m \geq 0$. If m is even, then c is a linear combination of b and 2; if m is odd, then c is even - in both cases we are done due to (1). $\square$

Therefore in the following we assume that $c > b > a > 2$.

Fact 2. Let $b > a > 2$ and suppose that $\gcd(a, b) = 1$. Then for $c \geq (a-1)(b-1)$ there exists a tame automorphism of $\mathbb{C}^3$ with multidegree (a, b, c) .

Proof. It is a classical fact due to Sylvester that every integer c greater than or equal to $c_0 = (a-1)(b-1)$ can be expressed as a linear combination of a and b with positive integer coefficients. Now apply Fact 1.(1). $\square$

Note that in this case $c_0 = \text{lcm}(a, b) - a - b + 1$. A similar reasoning is not valid if $\gcd(a, b) := d > 1$: since d divides every linear combination of a and b , we can only get that the automorphism with multidegree (a, b, c) exists for c divisible by d and $c \geq \text{lcm}(a, b) - a - b + d$. However, changing c_0 a little, we deal with the case $\gcd(a, b) > 1$ in the following theorem.

Theorem. Let $b > a > 2$ and set $r := \min\{b-1, (a-1)(\lfloor b/a \rfloor + 1)\}$. Then for $c \geq \text{lcm}(a, b) - r$ there exists a tame automorphism of $\mathbb{C}^3$ with multidegree equal to (a, b, c) .

Proof. Step 1. We will first show an automorphism for $c \geq \text{lcm}(a, b) - a$. Let $e := \text{lcm}(a, b)$ and write $c = e + (k-1)a + m$ for some $k \geq 0$ and $0 \leq m < a$. Let us take

$$\begin{aligned} F_1(x, y, z) &:= (x + z^a + z^m, y + z^b, z) \\ F_2(x, y, z) &:= (x, y, z + x^k(x^{e/a} - y^{e/b})) \end{aligned}$$

and consider the composition

$$F := F_2 \circ F_1 = (x + z^a + z^m, y + z^b, z + (x + z^a + z^m)^k v(x, y, z))$$

Then $v(x, y, z) = \sum_{i=1}^{e/a} \binom{e/a}{i} (x + z^m)^i z^{e-ia} - \sum_{j=1}^{e/b} \binom{e/b}{j} y^j z^{e-jb}$ and, provided that $m > 0$, the highest order monomial in v is $e/a z^m z^{e-a}$. Therefore for $m > 0$ the degree of the third coordinate of F equals $ka + m + e - a = c$. If $m = 0$, then $c = (e/a + k - 1)a$ and we can take $F = (x, y, z + x^{e/a+k-1}) \circ (x + z^a, y + z^b, z)$.

Step 2. Thanks to step 1, we must only find an automorphism F with multidegree (a, b, c) for $e - r \leq c < e - a$. Let us take $m := b + c - e$ (note that $0 < m < b$) and

$$\begin{aligned} F_1(x, y, z) &:= (x + z^a, y + z^b + z^m + u(x, z), z) \\ F_2(x, y, z) &:= (x, y, z + x^{e/a} - y^{e/b}) \end{aligned}$$

where u is a polynomial of degree $< b$. Consider the composition

$$F := F_2 \circ F_1 = (x + z^a, y + z^b + z^m + u(x, z), z + v_1(x, z) - v_2(x, y, z))$$

where $v_1(x, z) = \sum_{i=1}^{e/a} \binom{e/a}{i} x^i z^{e-ia}$, $v_2(x, y, z) = \sum_{j=1}^{e/b} \binom{e/b}{j} (y + z^m + u(x, z))^j z^{e-jb}$. Now let $u(x, z) = \sum_{k=1}^{\lfloor b/a \rfloor} u_k x^k z^{b-ka}$ and note that

$$u(x, z)^j = \sum_{s=j}^{j\lfloor b/a \rfloor} \sum_{k_1+\dots+k_j=s} u_{k_1} \cdot \dots \cdot u_{k_j} x^s z^{jb-sa}$$

where all the indices $k_1, \dots, k_j$ vary from 1 to $\lfloor b/a \rfloor$. One easily checks that the sum $\sum_{j=1}^{\lfloor b/a \rfloor} \binom{e/b}{j} u(x, z)^j z^{e-jb}$ contains all monomials of the form $x^i z^{e-ia}$ and degree higher than $e - b$ appearing in v_2 . Consequently, for $i \leq \lfloor b/a \rfloor$, coefficient of the term $x^i z^{e-ia}$ in $v_1 - v_2$ equals

$$\binom{e/a}{i} - \sum_{j=1}^{\lfloor b/a \rfloor} \binom{e/b}{j} \sum_{k_1+\dots+k_j=i} u_{k_1} \cdot \dots \cdot u_{k_j}$$

and becomes zero if $e/b u_i = \binom{e/a}{i} - \sum_{j=2}^{\lfloor b/a \rfloor} \binom{e/b}{j} \sum_{k_1+\dots+k_j=i} u_{k_1} \cdot \dots \cdot u_{k_j}$. Because on the right hand side the indices $k_1, \dots, k_j$ are smaller than i , the above formula allows a recursive definition of u_i , $1 \leq i \leq \lfloor b/a \rfloor$:

$$\begin{aligned} u_1 &:= b/a, \\ u_2 &:= b/e \left(\binom{e/a}{2} - \binom{e/b}{2} u_1^2 \right), \\ &\dots \\ u_i &:= b/e \left(\binom{e/a}{i} - \sum_{j=2}^{e/b} \binom{e/b}{j} \sum_{\substack{k_1+\dots+k_j=i \\ 1 \leq k_1, \dots, k_j \leq \lfloor b/a \rfloor}} u_{k_1} \cdot \dots \cdot u_{k_j} \right) \end{aligned}$$

Specifying u_i this way guarantees that the degree of the third coordinate of F is equal to the maximum of: $\deg(yx^{d-b})$, $\max\{\deg(x^i z^{d-ia}) : \lfloor b/a \rfloor < i \leq d/b\}$ and $\max\{\deg(y + z^m)^l u(x, z)^{j-l} z^{d-jb} : 1 \leq j \leq d/a, 1 \leq l \leq j\}$. An easy calculation shows that this maximum simplifies to $\max\{d - b + 1, d - (a - 1)(\lfloor b/a \rfloor + 1), m + d - b\} = \max\{d - r, c\} = c$. $\square$

Remark 1. If in the above theorem we get $r = b - 1$, then one can find an automorphism with multidegree (a, b, c) for $c \geq e - b$ (since $b|e - b$).

Definition 1. A pair (a, b) of positive integers such that $a < b$, $a \nmid b$ will be called tame iff for every $c > b$ there exists a tame automorphism of $\mathbb{C}^3$ with multidegree equal to (a, b, c) .

Corollary 1. A pair $(2n, kn)$, where $k \geq 3$ - odd is tame for $n \geq \frac{k-1}{2}$.

Proof. Due to Fact 1.(3) we can assume $n \geq 2$. We apply the Theorem and get the existence of tame automorphism with multidegree $(2n, kn, c)$, provided that $c \geq 2kn - r$, where $r = \min\{kn - 1, (2n - 1)(\lfloor kn/2n \rfloor + 1)\} = \min\{kn - 1, (2n - 1)(k + 1)/2\} = \min\{kn - 1, kn + n - \frac{k+1}{2}\}$. Following Remark 1, we are done if $n - \frac{k+1}{2} \geq -1$ i.e. $n \geq \frac{k-1}{2}$. $\square$

Corollary 2. Pairs $(2n, 3n)$ and $(2n, 5n)$ are tame for all $n \geq 1$.

Remark 2. In the case $n = \frac{k-3}{2}$ there are only following triples "left": $(4, 14, 15)$, $(6, 27, 28)$, $(8, 44, 45)$, ..., $(2n, (2n + 3)n, (2n + 3)n + 1)$.

REFERENCES

- [1] A. van den Essen, *Polynomial Automorphisms and the Jacobian Conjecture*, Progress in Math., vol. 190, Birkhäuser Verlag, Basel, Boston, Berlin, 2000.
- [2] M. Karaś, *There is no tame automorphism of $\mathbb{C}^3$ with multidegree $(3, 4, 5)$* , to appear.
- [3] W. van der Kulk, *On polynomial rings in two variables*, Nieuw Arch. Wisk. **1** (1953), pp. 34-41.
- [4] I. Shestakov, U. Umirbaev, *The tame and the wild automorphisms of polynomial rings in three variables*, J. Amer. Math. Soc. **17** (2004), pp. 197-227.

INSTITUTE OF MATHEMATICS, JAGIELLONIAN UNIVERSITY, ŁOJASIEWICZA 6, 30-348 KRAKÓW, POLAND.

E-mail address: jakub.zygadlo@im.uj.edu.pl